

The Two-Boson-Exchange Correction to Parity-Violating Elastic Electron-Proton Scattering

Shin Nan Yang¹, Keitaro Nagata², Hai Qing Zhou^{1,3}, Chung Wen Kao²

¹*Department of Physics and Center for Theoretical Sciences, National Taiwan University, Taipei 10617, Taiwan*

²*Department of Physics, Chung-Yuan Christian University, Chung-Li 32023, Taiwan*

³*Department of Physics, Southeast University, Nanjing, China*

Abstract

We calculate the two-boson-exchange (TBE) corrections to the parity-violating asymmetry of the elastic electron-proton scattering in a simple hadronic model including the nucleon and the $\Delta(1232)$ intermediate states. We find that Δ contribution δ_Δ is, in general, comparable with the nucleon contribution δ_N and the current experimental measurements of strange-quark effects in the proton neutral weak current. The total TBE corrections to the current extracted values of $G_E^s + \beta G_M^s$ in recent experiments are found to lie in the range of $-7 \sim +7\%$.

Key words: two-boson-exchange, parity-violation, elastic electron-proton scattering, strangeness form factor

PACS: 13.40.Ks, 13.60.Fz, 13.88.+e, 14.20.Dh

Strangeness content in the proton remains one of the most intriguing questions in hadron structure. The parity-violating asymmetry $A_{PV} = (\sigma_R - \sigma_L)/(\sigma_R + \sigma_L)$ in polarized electron elastic scattering provides a rather clean technique to probe the strangeness in the nucleon. Four experimental programs SAMPLE, HAPPEX, A4, and G0 [1] have been designed to measure this important quantity, which is small and ranges from 0.1 to 100 ppm. This calls for greater efforts to reduce theoretical uncertainty in order to arrive at a more reliable interpretation of experiments.

The radiative corrections to A_{PV} have been discussed in [2]. However, theoretical uncertainties remain. Recently, the contribution of the interference of the two-photon-exchange (TPE) process of Fig. 1(a) with Born diagrams has been evaluated in [3] in a parton model using GPDs. It was found that indeed the TPE correction to A_{PV} can reach several percent in certain kinematics, becoming comparable in size with existing experimental measurements of strange-quark effects in the proton neutral weak current. However, the partonic calculations of [3] are reliable only for Q^2 large comparable to a

typical hadronic scale, while all current experiments have been performed at lower Q^2 values. In addition, the γZ -exchange diagram of Fig. 1(b), expected to be of the same order as the TPE correction, was not considered in [3].

In this contribution, we report our recent calculations [4,5] of the TPE and γZ -exchange corrections to A_{PV} with N and $\Delta(1232)$ intermediate states as shown in Figs. 1(a) and 1(b), where both are treated in the same hadronic model developed in [6].

At hadron level, the couplings of photon and Z -boson with proton are well-known, while the couplings with the Δ are given as [5],




Fig. 1. (a) TPE, and (b) γZ -exchange diagrams for elastic electron-proton scattering. Corresponding cross-box diagrams are implied.

$$\begin{aligned}
\langle N(p') | J_\mu^\gamma | \Delta(p) \rangle &= \frac{F_\Delta(q^2)}{M_N^2} \bar{u}(p') [g_1(g_\mu^\alpha \not{p} \not{q} - p_\mu \gamma^\alpha \not{q} - \gamma_\mu \gamma^\alpha p \cdot q + \gamma_\mu \not{p} q^\alpha) \\
&\quad + g_2(p_\mu q^\alpha - p \cdot q g_\mu^\alpha) + g_3/M_N(q^2(p_\mu \gamma^\alpha - g_\mu^\alpha \not{p}) + q_\mu(q^\alpha \not{p} - \gamma^\alpha p \cdot q))] \gamma_5 T_3 u_\alpha^\Delta(p), \\
\langle p(p') | J_\mu^{Z,V} | \Delta^+(p) \rangle &= \frac{F_\Delta(q^2)}{M_N^2} \bar{u}(p') [\tilde{g}_1(g_\mu^\alpha \not{p} \not{q} - p_\mu \gamma^\alpha \not{q} - \gamma_\mu \gamma^\alpha p \cdot q + \gamma_\mu \not{p} q^\alpha) \\
&\quad + \tilde{g}_2(p_\mu q^\alpha - p \cdot q g_\mu^\alpha) + \tilde{g}_3/M_N(q^2(p_\mu \gamma^\alpha - g_\mu^\alpha \not{p}) + q_\mu(q^\alpha \not{p} - \gamma^\alpha p \cdot q))] \gamma_5 u_\alpha^\Delta(p), \\
\langle p(p') | J_\mu^{Z,A} | \Delta^+(p) \rangle &= \frac{H_\Delta(q^2)}{M_N^2} \bar{u}(p') [h_1(g_\mu^\alpha (p \cdot q) - p_\mu q^\alpha) + h_2/M_N^2(q^\alpha q_\mu \not{p} \not{q} \\
&\quad - (p \cdot q) \gamma^\alpha q_\mu \not{q}) + h_3((p \cdot q) \gamma^\alpha \gamma^\mu - \not{p} \gamma_\mu q^\alpha) + h_4(g_\mu^\alpha p^2 - p_\mu \gamma^\alpha \not{p})] u_\alpha^\Delta(p), \quad (1)
\end{aligned}$$

where M_N is the proton mass and $q = p' - p$, F 's and H 's are the corresponding form factors. The vertex functions in Eq. (1) satisfy constraints which ensure only the physical spin-3/2 component of the Rarita-Schwinger spinor couples with γ and Z [5].

The form factors F_Δ and H_Δ are chosen to have dipole form with a cut-off value of 0.84 and 1.0 GeV , respectively. $\tilde{g}_i$'s and g_i 's are related by $\tilde{g}_i = \sqrt{2/3} \alpha_V g_i$, with $\alpha_V = (1 - 2 \sin^2 \theta_W)/(2 \cos \theta_W)$. Coupling constants g_i 's are determined from $\gamma N \rightarrow \Delta$ at real photon point, while we infer the values of h_i 's from the data of $\nu N \rightarrow \mu \Delta$ [5].

We obtain the TPE and γZ -exchange corrections to A_{PV} as shown in Fig. 2, where we plot δ , defined by

$$A_{PV}(1\gamma + Z + 2\gamma + \gamma Z) = A_{PV}(1\gamma + Z)(1 + \delta_N + \delta_\Delta), \quad (2)$$

vs. $\epsilon \equiv [1 + 2(1 + \tau) \tan^2 \theta_{Lab}/2]^{-1}$, at two different values of $Q^2 = 0.1, 1.5 \text{ GeV}^2$. θ_{Lab} is the laboratory scattering angle and $\tau = Q^2/4M^2$. The subscripts N and Δ in δ refer to effect associated with N and Δ intermediate states, respectively.

We see from Fig. 2 that δ is sensitive *w.r.t.* both ϵ and Q^2 . For small Q^2 , δ_Δ is almost zero and negative at small ϵ , and decreases further as ϵ increases before turning around at $\epsilon \sim 0.8$. These behaviors are in sharp contrast with δ_N which always stays positive and decreases with increasing ϵ . Only at larger ϵ , δ_Δ is comparable with δ_N .

The strangeness form factors $G_E^s + \beta G_M^s$ extracted in the current experiments have already taken into account of the γZ -exchange effects as estimated in [2] within a zero-




Fig. 2. Two-boson-exchange corrections to parity-violating asymmetry as functions of ϵ from 0.1 to 0.99 at $Q^2 = 0.1, 1.5 \text{ GeV}^2$.

	I	II	III	IV	V	VI
$Q^2(\text{GeV}^2)$	0.477	0.109	0.23	0.108	0.262	0.410
ϵ	0.974	0.994	0.83	0.83	0.984	0.974
$\delta_N(\%)$	0.25	0.34	0.86	1.30	0.28	0.275
$\delta_\Delta(\%)$	1.20	1.90	-0.025	-0.65	1.33	1.14
$\delta_G(\%)$	+6.74	-7.53	-4.26	-4.82	-1.70	-0.10

Table 1

The corrections δ_G to $G_E^s + \beta G_M^s$ for HAPPEX, A4, and G0 experiments. (I, II), (III, IV), and (V, VI) refer to the HAPPEX, A4, and G0 data, respectively.

momentum-transfer approximation scheme. To avoid double counting, we need to first remove the γZ -exchange effects estimated with [2] and use our results instead. The resultant percentage change in values of the strange form factors defined as, $\overline{G}_E^s + \beta \overline{G}_M^s = (G_E^s + \beta G_M^s)(1 + \delta_G)$, where $\overline{G}_E^s + \beta \overline{G}_M^s$ denotes the values would be extracted with our estimation of TBE effects, is given in Table 1. We see that even though δ_N and δ_Δ are of the same sign and

of comparable size, the Δ effects actually reduce the difference between the results of [2] and the effects associated with only elastic nucleon intermediate states as found in [4]. The resultant corrections δ_G to the current extracted values for the strangeness form factor $G_E^s + \beta G_M^s$ are then only of a few percent. The large effects we find for both the nucleon and the Δ excitation brings up the question of the contributions of other higher resonances.

References

- [1] D.T. Spayde *et al.* (SAMPLE), Phys. Lett. **B 583**, 79 (2004); A. Acha *et al.* (HAPPEX), Phys. Rev. Lett. **98**, 032301 (2007); B. Glaser (for the A4 collaboration) Eur. Phys. J. **A 24**, S2, 141 (2005); C. Furget (for the G0 collaboration), Nucl. Phys. Proc. Suppl. **159**, 121 (2006); and references contained therein.
- [2] W.J. Marciano and A. Sirlin, Phys. Rev. **D 27**, 552 (1983); *ibid.* **29**, 75 (1984).
- [3] A.V. Afanasev, C.E. Carlson, Phys. Rev. Lett. **94**, 212301 (2005).
- [4] H.Q. Zhou, C.W. Kao, and S.N. Yang, Phys. Rev. Lett. **99**, 262001 (2007).
- [5] K. Nagata, H.Q. Zhou, C.W. Kao, and S.N. Yang, arXiv:0811.3539.
- [6] P.G. Blunden, W. Melnitchouk, and J.A. Tjon, Phys. Rev. Lett. **91**, 142304 (2003).